



# Séminaire Laurent Schwartz

## EDP et applications

Année 2014-2015


Patrick Gérard and Sandrine Grellier

**On the growth of Sobolev norms for the cubic Szegő equation**

*Séminaire Laurent Schwartz — EDP et applications* (2014-2015), Exposé n° XI, 20 p.

<[http://sisedp.cedram.org/item?id=SLSEDP\\_2014-2015\\_\\_\\_\\_A11\\_0](http://sisedp.cedram.org/item?id=SLSEDP_2014-2015____A11_0)>

© Institut des hautes études scientifiques & Centre de mathématiques Laurent Schwartz, École polytechnique, 2014-2015.

 Cet article est mis à disposition selon les termes de la licence  
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.  
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

Institut des hautes études scientifiques  
Le Bois-Marie • Route de Chartres  
F-91440 BURES-SUR-YVETTE  
<http://www.ihes.fr/>

Centre de mathématiques Laurent Schwartz  
UMR 7640 CNRS/École polytechnique  
F-91128 PALAISEAU CEDEX  
<http://www.math.polytechnique.fr/>

**cedram**

*Exposé mis en ligne dans le cadre du  
Centre de diffusion des revues académiques de mathématiques*  
<http://www.cedram.org/>

## ON THE GROWTH OF SOBOLEV NORMS FOR THE CUBIC SZEGŐ EQUATION

PATRICK GÉRARD AND SANDRINE GRELLIER

ABSTRACT. We report on a recent result establishing that trajectories of the cubic Szegő equation in Sobolev spaces with high regularity are generically unbounded, and moreover that, on solutions generated by suitable bounded subsets of initial data, every polynomial bound in time fails for high Sobolev norms. The proof relies on an instability phenomenon for a new nonlinear Fourier transform describing explicitly the solutions to the initial value problem, which is inherited from the Lax pair structure enjoyed by the equation.

### 1. INTRODUCTION

The large time behavior of solutions to Hamiltonian partial differential equations is an important problem in mathematical physics. In the case of finite dimensional Hamiltonian systems, many features of the large time behavior of trajectories are described using the topology of the phase space. For a given infinite dimensional system, several natural phase spaces, with different topologies, can be chosen, and the large time properties may strongly depend on the choice of such topologies. For instance, it is known that the cubic defocusing Schrödinger equation

$$i\partial_t u + \Delta u = |u|^2 u$$

posed on a Riemannian manifold  $M$  of dimension  $d = 1, 2, 3$  with sufficiently uniform properties at infinity, defines a global flow on the Sobolev spaces  $H^s(M)$  for every  $s \geq 1$  (see e.g. [4]). In this case, a typical large time behavior of interest is the boundedness of trajectories. On the energy space  $H^1(M)$ , the conservation of energy trivially implies that all the trajectories are bounded. On the other hand, the existence of unbounded trajectories in  $H^s(M)$  for  $s > 1$  is a long standing open problem [2], naturally connected to weak turbulence. In [3], [28], it was proved that dispersion properties imply a polynomial bound in time on  $H^s$  norms as time goes to infinity, but so far the optimality of this kind of bound has not been established or

---

2010 *Mathematics Subject Classification.* 35B15, 47B35, 37K15.

disproved.<sup>1</sup> Let us mention however that, for some special solutions of cubic NLS on  $\mathbb{R} \times \mathbb{T}^2$ , Hani–Pausader–Tzvetkov–Visciglia [17] succeeded in proving that  $H^s$  norms may not be bounded for  $s$  big enough, with a minoration of the type

$$e^{c(\log \log t)^{1/2}},$$

on a sequence of times going to infinity. Natural model problems for studying the unboundedness of Sobolev norms seem to be those for which the calculation of solutions is the most explicit, namely integrable systems. Typical examples are the Korteweg de Vries equation [21], [19] and the one dimensional cubic nonlinear Schrödinger defocusing equation [31], [15]. However, in these cases, the set of conservation laws is known to control the whole regularity of the solution, so that all the trajectories of elements of  $H^s(M)$  are bounded in  $H^s(M)$  for every nonnegative integer  $s$  [29].

In order to investigate direct and inverse cascades between spacial scales in one space dimension, Majda, Mc Laughlin, Tabak and their collaborators introduced, in a series of papers starting with [22], — see also Zakharov, Guyenne, Pushkarev, Dias [32] — a class of Hamiltonian equations on  $\mathbb{T}$  including

$$(1) \quad i\partial_t u - |D|^\alpha u = |u|^2 u, \quad D := \frac{1}{i} \partial_x, \quad \alpha > 0,$$

making numerical simulations suggesting weak turbulence effects for some values of  $\alpha$ . For  $\alpha > 1$ , it is easy to prove that this equation is globally well-posed on  $H^s(\mathbb{T})$  for  $s \geq \frac{\alpha}{2}$ , and that the  $H^{\alpha/2}$  norm is uniformly bounded along the trajectories. Moreover, polynomial bounds on the  $H^s$  norms for big  $s$  can also be obtained in that case [29].

In the limit case  $\alpha = 1$ , Equation (1) is a nonlinear (half-) wave equation in one space dimension, which can be proved to be globally well-posed on  $H^s$  for  $s \geq \frac{1}{2}$  (see [11]). However, compared to the case  $\alpha \neq 1$ , this equation is no more dispersive, which suggests that large time transition to high frequencies may be facilitated. In [11] — see also Pocovnicu [26]—, we proved that a Birkhoff normal form of this equation near the origin is given at first order by the following cubic Szegő equation,

$$(2) \quad i\partial_t u = \Pi(|u|^2 u),$$

where  $\Pi$  denotes the Szegő projector on  $L^2(\mathbb{T})$ ,

$$\Pi u(k) = \mathbf{1}_k \circ \hat{u}(k).$$

---

<sup>1</sup>Except in the case  $M = \mathbb{R}^d$ ,  $d = 2, 3, 4$ , where global finiteness of Strichartz norms has been established, leading to scattering theory [13], [27], [20], [6]. In this case, boundedness of high  $H^s$  norms follows. We are grateful to N. Tzvetkov for drawing our attention to this fact.

Notice that the range of  $\Pi$  is the closed subspace  $L_+^2(\mathbb{T})$  consisting of  $L^2$  functions on the unit circle which admit a holomorphic extension to the unit disc. Equation (2) has been introduced in [7], where global wellposedness was established on  $H_+^s(\mathbb{T}) := H^s(\mathbb{T}) \cap L_+^2(\mathbb{T})$  for every  $s \geq \frac{1}{2}$ , as well as boundedness of the  $H^{1/2}$  norm along every trajectory. Moreover, we proved that this equation enjoys a Lax pair structure, giving rise to integrability properties studied in [7] and [8], as well as the following a priori estimate on  $H^s$  norms of solutions,

$$(3) \quad \|u(t)\|_{H^s} \leq C_s e^{C_s t}, \quad s > 1,$$

where  $C_s$  is a uniform constant depending only on a bound of the  $H^s$  norm of the initial data. It turns out that this estimate is almost optimal, as shown by the following result.

**Theorem 1.**

- (1) For every  $s > \frac{1}{2}$ , the set of initial data in  $H_+^s(\mathbb{T})$  leading to an unbounded trajectory in  $H_+^s(\mathbb{T})$  is a dense  $G_\delta$ -subset of  $H_+^s(\mathbb{T})$ .
- (2) For every positive integer  $M$ , for every  $s > \frac{1}{2}$ , there exists a bounded subset  $B$  of  $\mathcal{C}_+(\mathbb{T}) := \cap_s H_+^s(\mathbb{T})$  such that

$$\sup_{u_0 \in B} \frac{\|u(t)\|_{H^s}}{(1+|t|)^M} \rightarrow \infty \text{ as } |t| \rightarrow \infty,$$

where  $u$  denotes the solution of (2) with  $u(0) = u_0$ .

The above theorem calls for several comments.

- It is in fact possible to find the same dense  $G_\delta$ -subset of initial data leading to unbounded trajectories in  $H^s$  for every  $s > \frac{1}{2}$ . However, at this stage we cannot display yet an explicit example of such an initial datum. For example, we know that every rational function with no pole in the closed unit disc gives rise to a bounded trajectory in every  $H^s$  (see [10]). This fact is in contrast with the case of the cubic Szegő equation on the line, where special examples of unbounded trajectories in  $H^s$ , for every  $s > \frac{1}{2}$ , are displayed in [25].
- In the above result, the superpolynomial growth of Sobolev norms is stated for families of solutions. In fact, we expect that this superpolynomial growth also holds for generic initial data, and that this growth is indeed exponential, making estimate (3) optimal. An explicit example of such a behaviour was recently displayed by H. Xu in [30], for the special perturbation

$$i\partial_t u = \Pi(|u|^2 u) + \alpha \int_{\mathbb{T}} u, \quad \alpha > 0.$$

- A very natural open question is of course whether Theorem 1 holds for the half-wave equation (1) for  $\alpha = 1$ .

The proof of Theorem 1 relies on explicit formulae for solutions of Equation (2), which are non trivial consequences of the Lax pair structure and will be presented without proof in section 2. Then we sketch the proof of part 1) of the theorem in section 3, and finally give a complete proof of part 2) in section 4.

## 2. EXPLICIT FORMULAE

In this section, we list the main formulae solving the Cauchy problem for the cubic Szegő equation (2). The detailed proofs will appear in a forthcoming paper.

First we introduce some additional notation. Given a positive integer  $n$ , we set

$$\Omega_n := \{s_1 > s_2 > \dots > s_n > 0\} \subset \mathbb{R}^n .$$

Given a nonnegative integer  $d \geq 0$ , we recall that a Blaschke product of degree  $d$  is a rational function on  $\mathbb{C}$  of the form

$$\Psi(z) = e^{-i\psi} \prod_{j=1}^d \frac{z - p_j}{1 - \bar{p}_j z} , \quad \psi \in \mathbb{T} , \quad p_j \in \mathbb{D} .$$

Alternatively,  $\Psi$  can be written as

$$\Psi(z) = e^{-i\psi} \frac{P(z)}{z^d \bar{P} \frac{1}{z}} ,$$

where  $\psi \in \mathbb{T}$  is called the angle of  $\Psi$  and  $P$  is a monic polynomial of degree  $d$  with all its roots in  $\mathbb{D}$ . Such polynomials are called Schur polynomials. We denote by  $\mathcal{B}_d$  the set of Blaschke products of degree  $d$ . It is a classical result — see *e.g.* [18] — that  $\mathcal{B}_d$  is diffeomorphic to  $\mathbb{T} \times \mathbb{R}^{2d}$ .

Given a  $n$ -tuple  $(d_1, \dots, d_n)$  of nonnegative integers, we set

$$\mathcal{S}_{d_1, \dots, d_n} := \Omega_n \times \prod_{r=1}^n \mathcal{B}_{d_r} ,$$

endowed with the natural product topology. Given a sequence  $(d_r)_{r=1}^n$  of nonnegative integers, we denote by  $\mathcal{S}_{(d_r)}^{(2)}$  the set of pairs

$$((s_r)_{r=1}^n, (\Psi_r)_{r=1}^n) \in \mathbb{R} \times \prod_{r=1}^n \mathcal{B}_{d_r}$$

such that

$$s_1 > \cdots > s_n > \cdots > 0, \quad (d_r + 1)s_r^2 < \infty, \quad r=1$$

We also endow  $\mathcal{S}_{(d_r)}^{(2)}$  with the natural topology.

Finally, we denote by  $\mathcal{S}_n$  the union of all  $\mathcal{S}_{d_1, \dots, d_n}$  over all the  $n$ -tuples  $(d_1, \dots, d_n)$ , and by  $\mathcal{S}^{(2)}$  the union of all  $\mathcal{S}_{(d_r)}^{(2)}$  over all the sequences  $(d_r)_{r=1}^\infty$ . Given  $(\mathbf{s}, \Psi) \in \mathcal{S}_n$  and  $z \in \mathbb{C}$ , we define the matrix  $\mathcal{C}(z) := \mathcal{C}(\mathbf{s}, \Psi)(z)$  as follows. If  $n = 2q$ , the coefficients of  $\mathcal{C}(\mathbf{s}, \Psi)(z)$  are given by

$$(4) \quad c_{jk}(z) := \frac{s_{2j-1} - s_{2k}z\Psi_{2k}(z)\Psi_{2j-1}(z)}{s_{2j-1}^2 - s_{2k}^2}, \quad j, k = 1, \dots, q.$$

If  $n = 2q - 1$ , we use the same formula as above, with  $s_{2q} = 0$ .

**Theorem 2.** *For every  $n \geq 1$ , for every  $(\mathbf{s}, \Psi) \in \mathcal{S}_n$ , for every  $z \in \overline{\mathbb{D}}$ , the matrix  $\mathcal{C}(\mathbf{s}, \Psi)(z)$  is invertible. We set*

$$(5) \quad u(\mathbf{s}, \Psi)(z) = \langle \mathcal{C}(z)^{-1}(\Psi_{2j-1}(z))_{1 \leq j \leq q}, \mathbf{1} \rangle,$$

where

$$\mathbf{1} := \begin{pmatrix} 1 \\ \cdot \\ \cdot \\ \cdot \\ 1 \end{pmatrix}, \quad \text{and } \langle X, Y \rangle := \sum_{k=1}^q X_k Y_k.$$

For every  $(\mathbf{s}, \Psi) \in \mathcal{S}^{(2)}$ , the sequence  $(u_q)_{q=1}^\infty$  with

$$u_q := u((s_1, \dots, s_{2q}), (\Psi_1, \dots, \Psi_{2q})),$$

is strongly convergent in  $H_+^{1/2}(\mathbb{S}^1)$ . We denote its limit by  $u(\mathbf{s}, \Psi)$ .

The mapping

$$(\mathbf{s}, \Psi) \in \bigcup_{n=1} \mathcal{S}_n \cup \mathcal{S}^{(2)} \longmapsto u(\mathbf{s}, \Psi) \in H_+^{1/2} \setminus \{0\}$$

is bijective. Furthermore, its restriction to every  $\mathcal{S}_{(d_1, \dots, d_n)}$  and to  $\mathcal{S}_{(d_r)}^{(2)}$  is a homeomorphism onto its range.

Finally, the solution at time  $t$  of equation (2) with initial data  $u_0 = u(\mathbf{s}, \Psi)$  is  $u(\mathbf{s}, \Psi)(t)$ , where

$$\Psi_r(t) = e^{i(-1)^r s_r^2 t} \Psi_r.$$

Let us briefly explain how the above nonlinear Fourier transform is related to spectral analysis. If  $u \in H_+^{1/2}(\mathbb{S}^1)$ , recall (see [23], [24]) that the Hankel operator of symbol  $u$  is the operator  $H_u : L_+^2(\mathbb{S}^1) \rightarrow L_+^2(\mathbb{S}^1)$  defined by

$$H_u(h) = \Pi(u\bar{h}) .$$

It can be shown that  $H_u^2$  is a positive selfadjoint trace class operator. If  $S$  is the shift operator defined by

$$Sh(z) = zh(z) ,$$

$H_u$  satisfies

$$S H_u = H_u S = H_{S u} .$$

We denote by  $K_u$  this new Hankel operator. Let us say that a positive real number  $s$  is a singular value associated to  $u$  if  $s^2$  is an eigenvalue of  $H_u^2$  or  $K_u^2$ . The main point in Theorem 2 is that the list  $s_1 > \dots > s_r > \dots$  is the list of singular values associated to  $u = u(\mathbf{s}, \Psi)$ , and that the corresponding list  $\Psi_1, \dots, \Psi_r, \dots$  describes the action of  $H_u$  and of  $K_u$  on the eigenspaces of  $H_u^2$ ,  $K_u^2$  respectively, making more precise a theorem of Adamyan-Arov-Krein about the structure of Schmidt pairs of Hankel operators [1]. In fact, denoting by  $u_j$  the orthogonal projection of  $u$  onto  $\ker(H_u^2 - s_{2j-1}^2 I)$  and by  $u_k$  the orthogonal projection of  $u$  onto  $\ker(K_u^2 - s_{2k}^2 I)$ , one can show that  $u_j \neq 0$ ,  $u_k \neq 0$ , and

$$\begin{aligned} \dim \ker(H_u^2 - s_{2j-1}^2 I) &= \deg \Psi_{2j-1} + 1, & \Psi_{2j-1}(z) H_u(u_j)(z) &= s_{2j-1} u_j(z), \\ \dim \ker(K_u^2 - s_{2k}^2 I) &= \deg \Psi_{2k} + 1, & K_u(u_k)(z) &= s_{2k} \Psi_{2k}(z) u_k(z) . \end{aligned}$$

**Remark 1.** These formulae have an interesting consequence in the special case where the Blaschke products  $\Psi_r$  are all equal to 1. Indeed, from formula (5), the Fourier coefficients of  $u$  are real, so that  $H_u$  and  $K_u$  act as self adjoint operators on the real subspace of  $L_+^2$  corresponding to real Fourier coefficients. Furthermore, from the above formulae, these operators are positive.

As a consequence of Theorem 2, we get inverse spectral theorems on Hankel operators, which generalize to singular values with arbitrary multiplicity the ones we had proved in [8] and [9] for simple singular values. Finally, the last assertion of Theorem 2 is a consequence of the following Lax pair identities,

$$\begin{aligned} \frac{dH_u}{dt} &= [B_u, H_u] , & B_u(h) &:= \frac{i}{2} H_u^2 h - i\Pi(|u|^2 h) , \\ \frac{dK_u}{dt} &= [C_u, K_u] , & C_u(h) &:= \frac{i}{2} K_u^2 h - i\Pi(|u|^2 h) . \end{aligned}$$

### 3. LARGE TIME INSTABILITY

Using Theorem 2, the proof of part 1) of Theorem 1 heavily relies on a new phenomenon, which is the loss of continuity of the map  $(\mathbf{s}, \Psi) \mapsto u(\mathbf{s}, \Psi)$  as three consecutive  $s_r$ 's are collapsing. More precisely, an inspection of formula (5) yields the following elementary lemma.

**Lemma 1.** *Given  $(s_1, \dots, s_n, \delta) \in \Omega_{n+1}$ ,  $(\Psi_1, \dots, \Psi_n) \in \mathcal{B}_{d_1} \times \dots \times \mathcal{B}_{d_n}$ , and  $\varepsilon > 0$  small enough, the family*

$$u^\varepsilon := u((s_1, \dots, s_n, \delta + \varepsilon, \delta, \delta - \varepsilon), (\Psi_1, \dots, \Psi_n, e^{i\varphi_+}, e^{i\theta}, e^{i\varphi_-}))$$

*converge as  $\varepsilon$  tends to 0, in every  $H^s$  space, to*

$$u = u((s_1, \dots, s_n, \delta), (\Psi_1, \dots, \Psi_n, \Psi))$$

*for some  $\Psi \in \mathcal{B}_1$ , except if  $\varphi_+ = \varphi_- \pmod{2\pi}$ . If  $\varphi_+ = \varphi_- \pmod{2\pi}$ , then  $u^\varepsilon$  is unbounded in  $H^s$  for every  $s > \frac{1}{2}$ .*

The mechanism of unboundedness of  $H^s$  norms for  $s > \frac{1}{2}$  in the case  $\varphi_+ = \varphi_- \pmod{2\pi}$  is the convergence, as  $\varepsilon$  tends to 0, of a pole of the rational function  $u^\varepsilon$  to the unit circle. Using this lemma, we get the following long time instability result.

**Proposition 1.** *For every  $u_0 \in C_+(\mathbb{T})$ , there exists a sequence  $(u_0^n)$  converging to  $u_0$  in  $C_+(\mathbb{T})$  and a sequence  $(t_n)$  of times such that the solution  $u^n$  of (2) with  $u^n(0) = u_0^n$  satisfies*

$$\forall s > \frac{1}{2}, \quad \|u^n(t_n)\|_{H^s} \xrightarrow{n} \infty.$$

Proposition 1 is a generalization to every smooth datum  $u_0$  of an instability phenomenon already displayed in [7], section 6, corollary 5, and revisited in [10], section 4, in the special case

$$u_0(z) = z.$$

This instability phenomenon was also discovered in [5] for the cubic NLS equation on  $\mathbb{T}^2$ , and made more precise, with polynomial estimates, in [14]. Indeed, we observed that, for every  $\varepsilon > 0$ , the solution  $u^\varepsilon$  of the cubic Szegő equation with the initial datum  $u_0^\varepsilon(z) = z + \varepsilon$  satisfies, at time  $t^\varepsilon \sim \frac{\pi}{2\varepsilon}$ ,

$$\forall s > \frac{1}{2}, \quad \|u^\varepsilon(t^\varepsilon)\|_{H^s} \simeq (t^\varepsilon)^{2s-1}.$$

This divergence is due to the existence of a pole of the rational function  $z \mapsto u^\varepsilon(t^\varepsilon, z)$  at distance  $d \simeq \varepsilon^2$  of the unit circle.

Let us sketch the proof of Proposition 1, which combines Lemma 1 with the last part of Theorem 2. By Theorem 2 and a standard density argument, we may assume that  $u_0$  has the form

$$u_0 = u((s_1, \dots, s_n), (\Psi_1, \dots, \Psi_n))$$



for some  $((s_1, \dots, s_n), (\Psi_1, \dots, \Psi_n)) \in \mathcal{S}_n$ . Given  $\delta > 0$ , Lemma 1 implies that

$$u_0^{\delta, \varepsilon} := u((s_1, \dots, s_n, \delta + \varepsilon, \delta, \delta - \varepsilon), (\Psi_1, \dots, \Psi_n, 1, 1, -1))$$

converges in every  $H^s$ , as  $\varepsilon$  tends to 0, to  $u((s_1, \dots, s_n, \delta), (\Psi_1, \dots, \Psi_n, \Psi))$  for some  $\Psi \in \mathcal{B}_1$ . Furthermore, it is easy to check that

$$u((s_1, \dots, s_n, \delta), (\Psi_1, \dots, \Psi_n, \Psi)) \longrightarrow u_0$$

in every  $H^s$  as  $\delta$  tends to 0. On the other hand, applying the last assertion of Theorem 2, the solution  $u^{\delta, \varepsilon}$  with the initial datum  $u_0^{\delta, \varepsilon}$  is given at time  $t$  by

$$u^{\delta, \varepsilon}(t) = u(((s_r)_{r=1}^n, \delta + \varepsilon, \delta, \delta - \varepsilon), ((e^{i(-1)^r t s_r^2} \Psi_r)_{r=1}^n, e^{i\varphi_+(t)}, e^{i\theta(t)}, e^{i\varphi_-(t)})),$$

with

$$\varphi_+(t) = (-1)^{n+1} t(\delta + \varepsilon)^2, \quad \theta(t) = (-1)^n t \delta^2, \quad \varphi_-(t) = \pi + (-1)^{n+1} t(\delta - \varepsilon)^2.$$

Choosing

$$t^{\delta, \varepsilon} = \frac{\pi}{4\delta\varepsilon},$$

we observe that the angles  $\varphi_+(t^{\delta, \varepsilon})$  and  $\varphi_-(t^{\delta, \varepsilon})$  are the same modulo  $2\pi$ , therefore, by Lemma 1,  $u^{\delta, \varepsilon}(t^{\delta, \varepsilon})$  is unbounded in  $H^s$  for every  $s > \frac{1}{2}$  as  $\varepsilon$  tends to 0. The proof of Proposition 1 then follows by choosing

$$u_0^n := u_0^{\delta_n, \varepsilon_n}, \quad t_n := t^{\delta_n, \varepsilon_n},$$

with appropriate sequences  $(\delta_n), (\varepsilon_n)$  satisfying  $\varepsilon_n \ll \delta_n \ll 1$ .

The first assertion of Theorem 1 is a consequence of Proposition 1 and of a Baire category argument. This argument comes back to Hani [16], who used it first for finding solutions with unbounded Sobolev norms for the totally resonant form of cubic NLS on  $\mathbb{T}^2$ , as investigated by [5] and [14]. Notice that the previously quoted result in [17] relies on [5] and [14], and on a remarkable modified scattering argument.

#### 4. THE FAILURE OF POLYNOMIAL ESTIMATES FOR HIGH SOBOLEV NORMS

In this section, we prove the super polynomial instability of the Sobolev norms for families of solutions of the Szegő cubic equation stated in the introduction. We recall this statement.

**Theorem 3.** *For every positive integer  $M$ , every  $s > \frac{1}{2}$ , there exists a bounded subset  $B$  of  $\mathcal{C}_+(\mathbb{S}^1)$  such that*

$$\sup_{u_0 \in B} \frac{\|u(t)\|_{H^s}}{(1 + |t|)^M} \rightarrow \infty \text{ as } |t| \rightarrow \infty$$

where  $u(t)$  is the solution of (2) with  $u(0) = u_0$ .

*Proof.* This result is a consequence of the following proposition.

**Proposition 2.** *For every  $N \geq 2$ , every  $\varepsilon > 0$ , every  $\xi := (\xi_j)_{1 \leq j \leq N} \in \mathbb{R}^N$ ,  $\eta := (\eta_k)_{1 \leq k \leq N-1} \in \mathbb{R}^{N-1}$ , such that*

$$\xi_1 > \eta_1 > \xi_2 > \eta_2 > \dots > \eta_{N-1} > \xi_N,$$

*we consider*

$$u_\varepsilon(z) = \langle \mathcal{C}_\varepsilon^{-1}(z)(\mathbf{1}), \mathbf{1} \rangle$$

*where  $\mathcal{C}_\varepsilon(z) = (c_{\varepsilon,jk}(z))_{1 \leq j,k \leq N}$ , with*

$$c_{\varepsilon,jk}(z) := \frac{1 + \varepsilon\xi_j - z(1 + \varepsilon\eta_k)}{(1 + \varepsilon\xi_j)^2 - (1 + \varepsilon\eta_k)^2}, \quad 1 \leq k \leq N-1;$$

$$c_{\varepsilon,jN}(z) := \frac{1}{1 + \varepsilon\xi_j}, \quad 1 \leq j \leq N.$$

*Then, there exists  $(\xi, \eta)$  such that*

- (1)  $\|u_\varepsilon\|_{H^s} \geq \frac{C}{\varepsilon^{(N-1)(2s-1)}}$ ;
- (2) *The family  $u_\varepsilon \frac{1}{2\varepsilon}, \cdot$  is bounded in  $\mathcal{C}_+(\mathbb{S}^1)$  as  $\varepsilon \rightarrow 0$ . Here  $(u_\varepsilon(t), \cdot)$  denotes the solution of (2) with  $u_\varepsilon(0) = u_\varepsilon$ .*

Theorem 3 is a direct consequence of this proposition by considering  $M = [(N-1)(2s-1)]$  and the family of initial data

$$B := u_\varepsilon \frac{1}{2\varepsilon}, \cdot, \quad \varepsilon > 0. \quad \square$$

The proof of proposition 2 requires several step. The first step consists in establishing that  $\|u_\varepsilon\|_{H^s} \geq \frac{C}{\varepsilon^{(N-1)(2s-1)}}$  for a dense open set of choices of  $(\xi, \eta)$ . This is based on the following lemma.

**Lemma 2.** *For a dense open set of  $(\xi, \eta)$ , we have  $(u_\varepsilon)(1) \geq \frac{C}{\varepsilon^{2(N-1)}}$ . Moreover, the poles of  $u_\varepsilon$  are simple,*

$$(6) \quad u_\varepsilon(z) = \alpha_0^\varepsilon + \sum_{j=1}^{N-1} \frac{\alpha_j^\varepsilon}{1 - p_j^\varepsilon z}$$

*with  $\alpha_0^\varepsilon, \alpha_j^\varepsilon > 0$ ,  $0 < p_j^\varepsilon < 1$ , and  $p_j^\varepsilon \rightarrow 1$  as  $\varepsilon \rightarrow 0$ . Furthermore,  $\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon)$  is equivalent to  $\varepsilon^{2(N-1)}$ .*

Let us first prove that the lemma implies that  $\|u_\varepsilon\|_{H^s} \geq \frac{C}{\varepsilon^{(N-1)(2s-1)}}$ . First, from (6) and the estimate on  $(u_\varepsilon)(1)$ ,

$$(u_\varepsilon)(1) = \sum_{j=1}^{N-1} \frac{\alpha_j^\varepsilon p_j^\varepsilon}{(1 - p_j^\varepsilon)^2} \geq \frac{C}{\varepsilon^{2(N-1)}}.$$

Hence, applying Cauchy-Schwarz, one gets

$$\begin{aligned}
 \frac{C}{\varepsilon^{2(N-1)}} &\leq \prod_{j=1}^{N-1} \frac{\alpha_j^\varepsilon p_j^\varepsilon}{(1-p_j^\varepsilon)^2} \\
 &\leq \prod_{j=1}^{N-1} \frac{(\alpha_j^\varepsilon)^2}{(1-p_j^\varepsilon)^{2s+1}} \quad 1/2 \quad \times \quad \prod_{j=1}^{N-1} \frac{(p_j^\varepsilon)^2}{(1-p_j^\varepsilon)^{3-2s}} \quad 1/2 \\
 &\leq \prod_{j=1}^{N-1} \frac{(\alpha_j^\varepsilon)^2}{(1-p_j^\varepsilon)^{2s+1}} \quad 1/2 \quad \times \quad \frac{C}{\varepsilon^{(3-2s)(N-1)}}.
 \end{aligned}$$

Eventually,

$$\prod_{j=1}^{N-1} \frac{(\alpha_j^\varepsilon)^2}{(1-p_j^\varepsilon)^{2s+1}} \quad 1/2 \quad \geq \frac{C}{\varepsilon^{(2s-1)(N-1)}}.$$

It remains to observe that

$$\|u_\varepsilon\|_{H^s}^2 \geq C \prod_{n=0}^{2s} n^{2s} \prod_{j=1}^{N-1} \alpha_j^\varepsilon (p_j^\varepsilon)^n \quad 2 \quad \geq C \prod_{j=1}^{N-1} \frac{(\alpha_j^\varepsilon)^2}{(1-p_j^\varepsilon)^{2s+1}}.$$

*Proof.* Let us turn to the proof of the lemma. We first estimate the derivative of  $u_\varepsilon$  at  $z = 1$ . From the explicit formula of  $u_\varepsilon$ , we obtain

$$(7) \quad (u_\varepsilon)'(1) = \langle \mathcal{C}_\varepsilon(1)^{-1} \dot{\mathcal{C}}_\varepsilon \mathcal{C}_\varepsilon(1)^{-1}(\mathbf{1}), \mathbf{1} \rangle = \langle \dot{\mathcal{C}}_\varepsilon \mathcal{C}_\varepsilon(1)^{-1}(\mathbf{1}), {}^t \mathcal{C}_\varepsilon(1)^{-1}(\mathbf{1}) \rangle$$

where  $\dot{\mathcal{C}}_\varepsilon = (\dot{c}_{\varepsilon,jk})_{1 \leq j,k \leq N}$ , with

$$\dot{c}_{\varepsilon,jk} := \frac{1 + \varepsilon \eta_k}{(1 + \varepsilon \xi_j)^2 - (1 + \varepsilon \eta_k)^2}, \quad 1 \leq k \leq N-1, \quad \dot{c}_{\varepsilon,jN} := 0, \quad 1 \leq j \leq N.$$

From the formula giving  $\mathcal{C}_\varepsilon$ , we have

$$\mathcal{C}_\varepsilon(1) = \left( \frac{1}{2 + \varepsilon(\xi_j + \eta_k)} \right)_{1 \leq j \leq N; 1 \leq k \leq N-1}, \quad \frac{1}{1 + \varepsilon \xi_j} \quad 1 \leq j \leq N$$

which is a Cauchy matrix. Let us recall that a Cauchy matrix is a matrix of the form  $\frac{1}{a_j + b_k}$ . Its determinant is given by

$$(8) \quad \det \frac{1}{a_j + b_k} = \frac{\prod_{i < j} (a_i - a_j) \prod_{k < l} (b_k - b_l)}{\prod_{j,k} (a_j + b_k)}.$$

It allows to obtain

$$\frac{1}{a_j + b_k} \quad -1 \quad = \quad (-1)^{j+k} \frac{\lambda_j \mu_k}{a_j + b_k} \quad kj$$

with

$$\lambda_j = \frac{l(a_j + b_l)}{i < j (a_i - a_j) \quad r > j (a_j - a_r)}$$

and

$$\mu_k = \frac{l(a_l + b_k)}{i < k (b_i - b_k) \quad r > k (b_k - b_r)}.$$

In the case of the matrix  $\mathcal{C}_\varepsilon(\mathbf{1})$ , we have

$$\lambda_j = \frac{2^{N-1} (1 + \varepsilon \xi_j)}{\varepsilon^{N-1}} \frac{l(1 + \varepsilon(\xi_j + \eta_l)/2)}{\xi_j}, \quad 1 \leq j \leq N,$$

and

$$\mu_k = \frac{2^N}{\varepsilon^{N-2}} \frac{l(1 + \varepsilon(\xi_l + \eta_k)/2)}{\eta_k(1 + \varepsilon \eta_k)}, \quad k \leq N - 1, \quad \mu_N = \frac{i(1 + \varepsilon \xi_i)}{k(1 + \varepsilon \eta_k)}$$

where we have set

$$\xi_j := \begin{matrix} (\xi_i - \xi_j) \\ i < j \end{matrix} \quad \begin{matrix} (\xi_j - \xi_r) \\ r > j \end{matrix}, \quad \eta_k := \begin{matrix} (\eta_i - \eta_k) \\ i < k \end{matrix} \quad \begin{matrix} (\eta_k - \eta_r) \\ r > k \end{matrix}.$$

Eventually, if  $k \leq N - 1$ ,

$$\begin{aligned} (\mathcal{C}_\varepsilon(\mathbf{1})^{-1}(\mathbf{1}))_k &= (-1)^k \mu_k \prod_{j=1}^N \frac{(-1)^j \lambda_j}{2 + \varepsilon(\xi_j + \eta_k)} \\ &= (-1)^k \mu_k \frac{2^{N-2}}{\varepsilon^{N-1}} \prod_{j=1}^N \frac{(-1)^j (1 + \varepsilon \xi_j)}{\xi_j} \prod_{l=k}^N \frac{1 + \varepsilon \frac{\xi_j + \eta_l}{2}}{2}; \end{aligned}$$

and

$$\begin{aligned} (\mathcal{C}_\varepsilon(\mathbf{1})^{-1}(\mathbf{1}))_N &= (-1)^N \mu_N \prod_{j=1}^N \frac{(-1)^j \lambda_j}{1 + \varepsilon \xi_j} \\ &= (-1)^N \mu_N \frac{2^{N-1}}{\varepsilon^{N-1}} \prod_{j=1}^N \frac{(-1)^j}{\xi_j} \prod_l \frac{1 + \varepsilon \frac{\xi_j + \eta_l}{2}}{2}. \end{aligned}$$

In order to simplify these quantities, we use the following identities.

**Lemma 3.** For any  $0 \leq p \leq N - 1$ ,

$$\prod_{j=1}^N (-1)^j \frac{\xi_j^p}{\xi_j} = \begin{cases} 0 & \text{if } p < N - 1 \\ -1 & \text{if } p = N - 1. \end{cases}$$

*Proof.* We view  $\prod_{j=1}^N (-1)^j \frac{\xi_j^p}{\xi_j}$  as a rational function of  $\xi_N$  denoted by  $Q_p(\xi_N)$ . Its poles are simple and equal to  $\xi_1, \dots, \xi_{N-1}$ . Identifying the residue at each of these poles, we get

$$\text{Res}(Q_p; \xi_N = \xi_r) = \frac{(-1)^{r+1} \xi_r^p}{\prod_{i < r} (\xi_i - \xi_r) \prod_{j > r} (\xi_r - \xi_j)} + \frac{(-1)^{N+1} \xi_r^p}{\prod_{i < N, i \neq r} (\xi_i - \xi_r)} = 0.$$

Hence  $Q_p(\xi_N)$  is in fact a polynomial. If  $p < N - 1$ , it tends to 0 as  $\xi_N$  tends to  $\infty$ , therefore it is identically 0. If  $p = N - 1$ , it is a constant, equal to its limit at infinity,

$$Q_p(\xi_N) = \frac{(-1)^N \xi_N^{N-1}}{(-1)^{N-1} \xi_N^{N-1}} = -1.$$

This completes the proof.  $\square$

Expanding in powers of  $\varepsilon$  the above formula giving  $\mathcal{C}_\varepsilon(\mathbf{1})^{-1}(\mathbf{1})_k$ , and using Lemma 3, we infer

$$(9) \quad \mathcal{C}_\varepsilon(\mathbf{1})^{-1}(\mathbf{1})_k = (-1)^{k+1} \mu_k, \quad k \leq N.$$

Let us compute  ${}^t \mathcal{C}_\varepsilon^{-1}(\mathbf{1})(\mathbf{1})$  in a similar way. From the formula, we have

$$\begin{aligned} {}^t \mathcal{C}_\varepsilon(\mathbf{1})^{-1}(\mathbf{1})_j &= (-1)^j \lambda_j \prod_{k=1}^{N-1} \frac{(-1)^k \mu_k}{2 + \varepsilon(\xi_j + \eta_k)} + \frac{(-1)^N \mu_N}{1 + \varepsilon \xi_j} \\ &=: (-1)^j \lambda_j \tilde{S}_j(\varepsilon). \end{aligned}$$

We expand in powers of  $\varepsilon$  and use again Lemma 3, changing  $N$  into  $N - 1$ . We get

$$\begin{aligned} \tilde{S}_j(\varepsilon) &= \frac{2^{N-1}}{\varepsilon^{N-2}} \prod_{k=1}^{N-1} \frac{(-1)^k}{\eta_k (1 + \varepsilon \eta_k)} \frac{1 + \varepsilon \frac{\xi_j + \eta_k}{2}}{2} + (-1)^N + O(\varepsilon) \\ &= (-1)^N + O(\varepsilon) \\ &\quad + \frac{2^{N-1}}{\varepsilon^{N-2}} \prod_{k=1}^{N-1} (-1)^k \varepsilon^{N-2} \prod_{p+q=N-2} (-1)^q C_{N-1}^p \frac{\eta_k^{p+q}}{2^p \eta_k} + O(\varepsilon^{N-1}) \\ &= -1 + O(\varepsilon). \end{aligned}$$

As a consequence, we infer

$$(10) \quad {}^t \mathcal{C}_\varepsilon(\mathbf{1})^{-1}(\mathbf{1})_j = (-1)^{j+1} \lambda_j (1 + O(\varepsilon)).$$

Eventually, inserting (9) and (10) into the formula (7) of  $(u_\varepsilon)$  (1), it gives

$$\begin{aligned} (u_\varepsilon) (1) &= \prod_{j=1}^N \prod_{k=1}^{N-1} \frac{1 + \varepsilon \eta_k}{(1 + \varepsilon \xi_j)^2 - (1 + \varepsilon \eta_k)^2} (-1)^{j+k} \lambda_j \mu_k (1 + O(\varepsilon)) \\ &= \frac{2^{2(N-1)}}{\varepsilon^{2(N-1)}} \prod_{j=1}^N \prod_{k=1}^{N-1} \frac{(-1)^{j+k}}{(\xi_j - \eta_k) \xi_j \eta_k} + O(\varepsilon) . \end{aligned}$$

Considering

$$\prod_{j,k=1}^{N-1} \frac{(-1)^{j+k}}{(\xi_j - \eta_k) \xi_j \eta_k}$$

as a function of  $\xi_N$ , the pole  $\eta_{N-1}$  appears only once. Hence, this meromorphic function does not vanish on an open dense set of  $(\xi, \eta)$ . This proves that

$$|(u_\varepsilon) (1)| \geq \frac{C}{\varepsilon^{2(N-1)}}$$

for an open dense set of choices of  $(\xi, \eta)$ .

Next we prove the second statement of Lemma 2. Thanks to remark 1, operators  $H_{u_\varepsilon}$  and  $K_{u_\varepsilon}$  act as positive self adjoint operators on the subspace of  $L_+^2$  corresponding to real Fourier coefficients. As a consequence — see e.g. [12], Prop. 2.1 —, there exists a bounded positive measure  $\mu$  on  $[0, 1[$  such that

$$\forall k \geq 0 , \hat{u}_\varepsilon(k) = \int_{[0,1[} t^k d\mu(t) .$$

On the other hand,  $H_{u_\varepsilon}$  has exactly  $N$  positive singular values, hence is an operator of finite rank  $N$ . By the identity  $S H_{u_\varepsilon} = H_{u_\varepsilon} S$ ,  $S$  acts on the range of  $H_{u_\varepsilon}$ , therefore there exists a non trivial linear relation between the vectors  $(S)^p u_\varepsilon$ ,  $p = 0, \dots, N$ . Equivalently, there exists non trivial coefficients  $c_p$  such that

$$\forall k \geq 0 , \sum_{p=0}^N c_p \hat{u}(k+p) = 0 .$$

This precisely means that there exists a non trivial polynomial  $P$  of degree  $N$  such that

$$\forall k \geq 0 , \int_{[0,1[} t^k P(t) d\mu(t) = 0 .$$

By the Weierstrass theorem,  $\mu$  is therefore a positive linear combination of Dirac measures. This implies that the poles of  $u_\varepsilon$  are simple and that

$$u_\varepsilon(z) = \alpha_0^\varepsilon + \sum_{j=1}^{N-1} \frac{\alpha_j^\varepsilon}{1 - p_j^\varepsilon z}$$

with  $\alpha_j^\varepsilon > 0$ ,  $0 < p_j^\varepsilon < 1$ . In particular  $(u_\varepsilon)(1) > 0$  so that

$$(u_\varepsilon)(1) \geq \frac{C}{\varepsilon^{2(N-1)}}.$$

Let us show that  $p_j^\varepsilon \rightarrow 1$  as  $\varepsilon \rightarrow 0$ . We know that the denominator of  $u_\varepsilon(z)$  is

$$P_\varepsilon(z) := \frac{\det \mathcal{C}_\varepsilon(z)}{\det \mathcal{C}_\varepsilon(0)} = \prod_{j=1}^{N-1} (1 - p_j^\varepsilon z),$$

therefore it suffices to prove that  $P_\varepsilon(z) \simeq (1 - z)^{N-1}$  as  $\varepsilon$  tends to 0. From the formula of  $\mathcal{C}_\varepsilon(z)$ ,

$$(2\varepsilon)^{N-1} \det \mathcal{C}_\varepsilon(z) \rightarrow (1 - z)^{N-1} \det \begin{pmatrix} 1 & & & \\ & \xi_j - \eta_k & & \\ & & \ddots & \\ & & & \xi_{j-1} - \eta_{k-1} \end{pmatrix}, \mathbf{1}$$

and

$$(2\varepsilon)^{N-1} \det \mathcal{C}_\varepsilon(0) \rightarrow \det \begin{pmatrix} 1 & & & \\ & \xi_j - \eta_k & & \\ & & \ddots & \\ & & & \xi_{j-1} - \eta_{k-1} \end{pmatrix}, \mathbf{1},$$

so the claim is proved.

Finally, we establish that

$$\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon) \simeq \varepsilon^{2(N-1)}.$$

Let us first prove that  $\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon) \leq C\varepsilon^{2(N-1)}$ . From the estimate of  $(u_\varepsilon)(1)$ ,

$$\begin{aligned} \frac{C}{\varepsilon^{2N-1}} &\leq (u_\varepsilon)(1) = \frac{\alpha_j^\varepsilon p_j^\varepsilon}{(1 - p_j^\varepsilon)^2} \\ &\leq \frac{1}{\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon)} \frac{\alpha_j^\varepsilon p_j^\varepsilon}{(1 - p_j^\varepsilon)} \\ &\leq \frac{u_\varepsilon(1)}{\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon)} \leq \frac{\text{tr}(H_{u_\varepsilon}) + \text{tr}(K_{u_\varepsilon})}{\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon)} \\ &\leq \frac{2N - 1 + O(\varepsilon)}{\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon)}. \end{aligned}$$

It gives the bound from above for  $\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon)$ . For the bound from below, we use formula (8) to obtain

$$\begin{aligned} (1 - p_j^\varepsilon) &= \frac{\det \mathcal{C}_\varepsilon(1)}{\det \mathcal{C}_\varepsilon(0)} \simeq \varepsilon^{N-1} \det \mathcal{C}_\varepsilon(1) \\ &\simeq \varepsilon^{N-1} \det \begin{pmatrix} 1 & & & \\ 2 + \varepsilon(\xi_j + \eta_k) & & & \\ & & \ddots & \\ & & & 2 + \varepsilon(\xi_{j-1} + \eta_{k-1}) \end{pmatrix}, \frac{1}{1 + \varepsilon \xi_j} \\ &\simeq \varepsilon^{N(N-1)}. \end{aligned}$$

On the other hand, as

$$-P_\varepsilon(1) = \prod_{j=1}^N p_j^\varepsilon \prod_{i=j}^N (1 - p_i^\varepsilon) \simeq \frac{(1 - p_1^\varepsilon)}{\min_{1 \leq j \leq N} (1 - p_j^\varepsilon)} \simeq \frac{\varepsilon^{N(N-1)}}{\min_{1 \leq j \leq N} (1 - p_j^\varepsilon)},$$

it suffices to bound  $-P_\varepsilon(1)$  from above. Using formula (8) again,

$$\begin{aligned} -\frac{d}{dz}(\det \mathcal{C}_\varepsilon(z))|_{z=1} &= \prod_{k=1}^{N-1} \prod_{l=1}^N \frac{(-1)^{k+l}(1 + \varepsilon\eta_k)}{(1 + \varepsilon\xi_j)^2 - (1 + \varepsilon\eta_k)^2} \\ &\quad \cdot \det \frac{1}{2 + \varepsilon(\xi_j + \eta_m)}_{j=l, m=k}, \quad \frac{1}{1 + \varepsilon\xi_j}_{j=l} \\ &\leq \varepsilon^{(N-2)^2-1} \end{aligned}$$

Eventually,

$$-P_\varepsilon(1) \simeq -\varepsilon^{N-1} \frac{d}{dz}(\det \mathcal{C}_\varepsilon(z))|_{z=1} \leq C\varepsilon^{(N-1)(N-2)}$$

so that

$$\min_{1 \leq j \leq N-1} (1 - p_j^\varepsilon) \geq C\varepsilon^{2(N-1)}.$$

This completes the proof of part 1 of Proposition 2.  $\square$

We are left with proving the second part of Proposition 2. From Theorem 2, at time  $t = \frac{1}{2\varepsilon}$

$$u_\varepsilon = \frac{1}{2\varepsilon}, z = \langle \tilde{\mathcal{C}}_\varepsilon(z)^{-1}(\Psi_{\text{odd}}^\varepsilon), \mathbf{1} \rangle$$

where  $\Psi_{\text{odd}}^\varepsilon := (e^{-i\psi_j^\varepsilon(1/2\varepsilon)})_{1 \leq j \leq N}$  with  $\psi_j^\varepsilon(t) := (1 + \varepsilon\xi_j)^2 t$ . Hence

$$\Psi_{\text{odd}}^\varepsilon = e^{-i/2\varepsilon}(1 + O(\varepsilon)) e^{-i\xi_j}_{1 \leq j \leq N},$$

and, for  $1 \leq j \leq N$ ,

$$\begin{aligned} \tilde{\mathcal{C}}_\varepsilon(z)_{jk} &= \frac{1 + \varepsilon\xi_j - z(1 + \varepsilon\eta_k)e^{-i(\xi_j - \eta_k)}(1 + O(\varepsilon))}{(1 + \varepsilon\xi_j)^2 - (1 + \varepsilon\eta_k)^2}, \quad 1 \leq k \leq N-1, \\ \tilde{\mathcal{C}}_\varepsilon(z)_{jN} &= \frac{1}{1 + \varepsilon\xi_j}. \end{aligned}$$

Let  $\tilde{P}_\varepsilon(z) = \frac{\det \tilde{\mathcal{C}}_\varepsilon(z)}{\det \tilde{\mathcal{C}}_\varepsilon(0)}$ . From Theorem 2,  $\tilde{P}_\varepsilon(z)$  has  $(N-1)$  roots outside the closed unit disc  $\overline{\mathbb{D}}$ . Our aim is to prove that, for a suitable choice of  $(\xi, \eta)$ , the distance of these roots to the boundary of  $\overline{\mathbb{D}}$  is bounded from below. Define

$$P(\xi, \eta)(z) := \lim_{\varepsilon \rightarrow 0} \tilde{P}_\varepsilon(z).$$



Observe that

$$\begin{aligned} P(\xi, \eta)(z) &= c(\xi, \eta) \lim_{\varepsilon \rightarrow 0} \varepsilon^{N-1} \det \tilde{\mathcal{C}}_\varepsilon(z) \\ &= \tilde{c}(\xi, \eta) \det \begin{pmatrix} 1 - ze^{-i(\xi_j - \eta_k)} \\ \xi_j - \eta_k \end{pmatrix}_{1 \leq j \leq N-1, 1 \leq k \leq N-1}, \mathbf{1} \end{aligned}$$

Notice that the determinant in the right hand side is a polynomial in  $z$  of degree  $N - 1$  whose coefficient of  $z^{N-1}$  equals

$$(-1)^{N-1} \det \begin{pmatrix} e^{-i\xi_j} \\ \xi_j - \eta_k \end{pmatrix}_{1 \leq j \leq N-1, 1 \leq k \leq N-1}, \mathbf{1}.$$

With the choice  $\xi = \xi = (2\pi(N - j + 1))_{1 \leq j \leq N}$ , this determinant is

$$(-1)^{N-1} \det \begin{pmatrix} e^{i\eta_k} \\ \xi_j - \eta_k \end{pmatrix}_{1 \leq j \leq N-1, 1 \leq k \leq N-1}, \mathbf{1} \neq 0.$$

The non vanishing of this determinant follows from developing with respect to the last column, and in view of the explicit formula for the Cauchy determinants. By continuity, this determinant remains non zero in a neighborhood of  $\xi = \xi$ . Furthermore, Theorem 2 tells us that the roots of  $\tilde{P}_\varepsilon(z)$  are located outside the unit disc. Hence all the roots of  $P(\xi, \eta)$  belong to  $\{z, |z| \geq 1\}$ . Furthermore,

$$P(\xi, \eta)(z) = \tilde{c}(\xi, \eta) \det \begin{pmatrix} 1 \\ \xi_j - \eta_k \end{pmatrix}_{1 \leq j \leq N-1, 1 \leq k \leq N-1}, \mathbf{1} \prod_{k=1}^{N-1} (1 - ze^{i\eta_k}).$$

Hence, for a suitable choice of  $\eta$  so that  $\eta_k \neq \eta_l + 2\pi$  for  $k \neq l$ ,  $P(\xi, \eta)$  has only simple zeroes which belong to the unit circle.

**Lemma 4.** *For any  $\eta$  such that  $P(\xi, \eta)$  has only simple zeroes on the unit circle, there exists an open neighborhood of  $\xi$  such that, for every  $\xi \neq \xi$  in this neighborhood, the zeroes of  $P(\xi, \eta)$  are all outside  $\mathbb{D}$ .*

*Proof.* Denote by

$$\{z_k(\xi, \eta); 1 \leq k \leq N - 1\}$$

the simple zeroes of  $P(\xi, \eta)$ , with  $z_k(\xi, \eta) = e^{-i\eta_k}$ . The functions  $\xi \mapsto z_k(\xi, \eta)$  are analytic and satisfy  $|z_k(\xi, \eta)|^2 \geq 1$  and  $|z_k(\xi, \eta)|^2 = 1$ . In particular the quadratic form  $\xi \mapsto Q_k(\xi)$  associated to the Hessian of the function  $\xi \mapsto |z_k(\xi, \eta)|^2$  is positive at any  $\xi$ . We want to prove that, for any  $k$ ,  $Q_k$  is not identically 0 for  $\eta$  in a dense open set. It suffices to prove that the Laplacian of  $\xi \mapsto |z_k(\xi, \eta)|^2$ , which coincides with the trace of  $Q_k$ , is not identically zero. Let us compute this Laplacian.

$$\sum_{j=1}^N \frac{\partial^2}{\partial \xi_j^2} \frac{|z_k|^2}{2} \Big|_{\xi=\xi} = \sum_{j=1}^N \operatorname{Re} \bar{z}_k \frac{\partial^2 z_k}{\partial \xi_j^2} \Big|_{\xi=\xi} + \sum_{j=1}^N \frac{\partial z_k^2}{\partial \xi_j} \Big|_{\xi=\xi}.$$

Differentiating the equation  $P(\xi, \eta)(z_k(\xi)) = 0$ , we obtain

$$\frac{\partial z_k}{\partial \xi_j} = -\frac{\frac{\partial P}{\partial \xi_j}}{\frac{\partial P}{\partial z}}$$

and

$$\frac{\partial^2 z_k}{\partial \xi_j^2} = \frac{-\frac{\partial^2 P}{\partial \xi_j^2}}{\frac{\partial P}{\partial z}} + 2 \frac{\frac{\partial^2 P}{\partial \xi_j \partial z} \frac{\partial P}{\partial \xi_j}}{\frac{\partial P}{\partial z}^2} - \frac{\frac{\partial P}{\partial \xi_j}^2 \frac{\partial^2 P}{\partial z^2}}{\frac{\partial P}{\partial z}^3}$$

Introduce the following quantities.

$$D_{jk} := \frac{1}{\xi_j - \eta_k} \det \begin{matrix} 1 \\ \xi_r - \eta_l \end{matrix}_{r=j, l=k}, \mathbf{1},$$

$$D := \prod_j (-1)^{j+k} D_{jk}, \zeta_k := \prod_{l=k} (1 - e^{i(\eta_l - \eta_k)})$$

$$a_{lk} := -\frac{e^{i(\eta_l - \eta_k)}}{1 - e^{i(\eta_l - \eta_k)}} = \frac{e^{i(\eta_l - \eta_k)/2}}{2i \sin(\eta_l - \eta_k)/2}.$$

Notice that  $\text{Re}(a_{lk}) = \frac{1}{2}$ . Differentiating

$$P(\xi, \eta)(z) = \det \begin{matrix} 1 - ze^{-i(\xi_j - \eta_k)} \\ \xi_j - \eta_k \end{matrix}_{1 \leq j \leq N; 1 \leq k \leq N-1}, \mathbf{1},$$

one gets, after some computations,

$$\frac{\partial P}{\partial \xi_j} \Big|_{\xi=\xi, z=z_k} = i(-1)^{j+k} D_{jk} \zeta_k, \quad \frac{\partial P}{\partial z} \Big|_{\xi=\xi, z=z_k} = -e^{i\eta_k} \zeta_k D$$

$$\frac{\partial^2 P}{\partial \xi_j^2} \Big|_{\xi=\xi, z=z_k} = (-1)^{j+k} \zeta_k D_{jk} \left( 1 - \frac{2i}{\xi_j - \eta_k} \right)$$

$$\frac{\partial^2 P}{\partial z^2} \Big|_{\xi=\xi, z=z_k} = 2 \sum_{l=k} \frac{\zeta_k e^{i(\eta_l + \eta_k)}}{1 - e^{i(\eta_l - \eta_k)}} D = -2 \sum_{l=k} \zeta_k e^{2i\eta_k} a_{lk} D,$$

and

$$\begin{aligned} \frac{\partial^2 P}{\partial \xi_j \partial z} \Big|_{\xi=\xi, z=z_k} &= (-1)^{j+k} \zeta_k D_{jk} e^{i\eta_k} \left( i + \frac{1}{\xi_j - \eta_k} - i \sum_{l=k} \frac{e^{i(\eta_l - \eta_k)}}{1 - e^{i(\eta_l - \eta_k)}} \right) \\ &\quad + \zeta_k e^{i\eta_k} \sum_{l=k} (-1)^{j+l} D_{jl} \left( \frac{1}{\xi_j - \eta_l} - i \frac{e^{i(\eta_l - \eta_k)}}{1 - e^{i(\eta_l - \eta_k)}} \right) \\ &= \zeta_k e^{i\eta_k} (-1)^{j+k} D_{jk} \left( i + \frac{1}{\xi_j - \eta_k} + i \sum_{l=k} a_{lk} \right) \\ &\quad + \sum_{l=k} (-1)^{j+l} D_{jl} \left( \frac{1}{\xi_j - \eta_l} + i a_{lk} \right). \end{aligned}$$

Hence, inserting these formulae to compute the Laplacian, we obtain

$$\begin{aligned} \operatorname{Re} \bar{z}_k \frac{\partial^2 z_k}{\partial \xi_j^2} \Big|_{\xi=\xi} &= \operatorname{Re} e^{i\eta_k} \frac{\partial^2 z_k}{\partial \xi_j^2} \Big|_{\xi=\xi} \\ &= I_j + II_j + III_j \end{aligned}$$

with

$$\begin{aligned} I_j &= \operatorname{Re} e^{i\eta_k} \frac{-\frac{\partial^2 P}{\partial \xi_j^2}}{\frac{\partial P}{\partial z}} = (-1)^{j+k} \frac{D_{jk}}{D}, \\ II_j &= 2 \operatorname{Re} e^{i\eta_k} \frac{\frac{\partial^2 P}{\partial \xi_j \partial z} \frac{\partial P}{\partial \xi_j}}{\frac{\partial P}{\partial z}} \\ &= 2 \operatorname{Re} \frac{i D_{jk}^2 \left( i + \frac{1}{\xi_j - \eta_k} + i \sum_{l=k} a_{lk} \right) + \sum_{l=k} (-1)^{k+l} D_{jk} D_{jl} \frac{1}{\xi_j - \eta_l} + i a_{lk}}{D^2} \\ &= \frac{D_{jk}^2 \left( -2 - (N-2) - \sum_{l=k} (-1)^{k+l} D_{jk} D_{jl} \right)}{D^2}, \\ III_j &= -\operatorname{Re} e^{i\eta_k} \frac{\frac{\partial P}{\partial \xi_j} \frac{\partial^2 P}{\partial z^2}}{\frac{\partial P}{\partial z}} = \operatorname{Re} \frac{2 \sum_{l=k} a_{lk} D_{jk}^2}{D^2} = (N-2) \frac{D_{jk}^2}{D^2}. \end{aligned}$$

Remark that

$$\sum_{l=k} (-1)^{j+l} D_{jl} = D - (-1)^{j+N} \det \frac{1}{\xi_r - \eta_l} \Big|_{r=j}.$$

Hence, it gives

$$\begin{aligned} \sum_{j=1}^N \frac{\partial^2}{\partial \xi_j^2} \frac{|z_k|^2}{2} \Big|_{\xi=\xi} &= \sum_{j=1}^N \sum_l (-1)^{k+l+1} \frac{D_{jk} D_{jl}}{D^2} + (-1)^{j+k} \frac{D_{jk}}{D} \\ &= (-1)^{k+N} \frac{1}{D^2} \sum_j D_{jk} \det \frac{1}{\xi_r - \eta_l} \Big|_{r=j} \end{aligned}$$

It remains to check that

$$(-1)^{k+N} \sum_j D_{jk} \det \frac{1}{\xi_r - \eta_l} \Big|_{r=j}$$

is not identically zero. This follows from the fact that the limit as  $\eta_k$  tends to infinity of

$$(-1)^{k+N} \eta_k^2 \sum_j D_{jk} \det \frac{1}{\xi_r - \eta_l} \Big|_{r=j}$$

equals

$$\det_j \frac{1}{\xi_r - \eta_l} \Big|_{r=j, l=k}^2, \mathbf{1}$$

which is clearly not identically zero for a dense choice of  $(\eta, \xi)$ .  $\square$

## REFERENCES

- [1] Adamyan, V. M., Arov, D. Z., Krein, M. G., *Analytic properties of the Schmidt pairs of a Hankel operator and the generalized Schur-Takagi problem.* (Russian) Mat. Sb. (N.S.) 86(128) (1971), 34–75; English transl. Math USSR. Sb. 15 (1971), 31–73.
- [2] Bourgain, J., *Problems in Hamiltonian PDE's*, Geom. Funct. Anal. (2000), Special Volume, Part I, 32–56.
- [3] Bourgain, J., *On the growth in time of higher Sobolev norms of smooth solutions of Hamiltonian PDE*, Internat. Math. Res. Notices, (1996), 277–304.
- [4] Burq, N., Gérard, P., Tzvetkov, N.: *Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds* Amer. J. Math. 126, 569–605 (2004).
- [5] Colliander J., Keel M., Staffilani G., Takaoka H., Tao, T., *Transfer of energy to high frequencies in the cubic defocusing nonlinear Schrödinger equation*, Inventiones Math. 181 (2010), 39–113.
- [6] Dodson, B., *Global well-posedness and scattering for the defocusing,  $L^2$ -critical, nonlinear Schrödinger equation when  $d = 2$* , arXiv:1006.1375, preprint, 2011.
- [7] Gérard, P., Grellier, S., *The cubic Szegő equation*, Ann. Scient. Éc. Norm. Sup. 43 (2010), 761–810.
- [8] Gérard, P., Grellier, S., *Invariant Tori for the cubic Szegő equation*, Invent. Math. 187 (2012), 707–754.
- [9] Gérard, P., Grellier, S., *Inverse spectral problems for compact Hankel operators*, J. Inst. Math. Jussieu 13 (2014), 273–301.
- [10] Gérard, P., Grellier, S., *An explicit formula for the cubic Szegő equation*, to appear in Trans. A.M.S.
- [11] Gérard, P., Grellier, S., *Effective integrable dynamics for a certain nonlinear wave equation*, Anal. PDEs 5 (2012), 1139–1155.
- [12] Gérard, P., Pushnitski, A., *An inverse problem for self-adjoint positive Hankel operators*, arXiv:1401.2042, to appear in IMRN.
- [13] Ginibre, J., Velo, G. *Scattering theory in the energy space for a class of nonlinear Schrödinger equations*, J. Math. Pures Appl. (9) 64 (1985), no. 4, 363–401.
- [14] Guardia, M., Kaloshin, V., *Growth of Sobolev norms in the cubic defocusing nonlinear Schrödinger equation*, arXiv:1205.5188[math.AP], to appear in J. European Math. Soc.
- [15] Grébert, B., Kappeler, T., *The defocusing NLS equation and Its Normal Form*, EMS series of Lectures in Mathematics, European Mathematical Society, 2014.
- [16] Hani, Z., *Long-time strong instability and unbounded orbits for some periodic nonlinear Schrödinger equations*, to appear in Archives for Rational Mechanics and Analysis. arXiv:1210.7509.
- [17] Hani, Z., Pausader, B., Tzvetkov, N., Visciglia, N., *Modified scattering for the cubic Schrödinger equations on product spaces and applications*, arXiv:1311.2275, 2013.

- [18] Hernández B. A., Frías-Armenta M. E., Verduzco F., *On differential structures of polynomial spaces in control theory*, Journal of Systems Science and Systems Engineering 21 (2012), 372–382.
- [19] Kappeler, T., Pöschel, J., *KdV & KAM*, A Series of Modern Surveys in Mathematics, vol. 45, Springer-Verlag, 2003.
- [20] Killip, R., Tao, T., Visan, M., *The cubic nonlinear Schrödinger equation in two dimensions with radial data*. J. Eur. Math. Soc. 11 (2009), 1203–1258.
- [21] Lax, P. : *Integrals of Nonlinear equations of Evolution and Solitary Waves*, Comm. Pure and Applied Math. 21, 467–490 (1968).
- [22] Majda, A., Mc Laughlin, D., Tabak, E., *A one dimensional model for dispersive wave turbulence*, J. Nonlinear Sci. 7 (1997) 9–44.
- [23] Nikolskii, N. K., *Operators, functions, and systems: an easy reading. Vol. 1. Hardy, Hankel, and Toeplitz*. Translated from the French by Andreas Hartmann. Mathematical Surveys and Monographs, 92. American Mathematical Society, Providence, RI, 2002.
- [24] Peller, V.V., *Hankel Operators and their applications* Springer Monographs in Mathematics. Springer-Verlag, New York, 2003.
- [25] Pocovnicu, O. *Explicit formula for the solution of the Szegő equation on the real line and applications*, Discrete Cont. Dyn. Syst. 31 (2011), 607–649.
- [26] Pocovnicu, O. *First and second order approximations of a nonlinear wave equation* J. Dynam. Differential Equations, article no. 9286 (2013), 29 pp, DOI:10.1007/s10884-013-9286-5.
- [27] Ryckman, E., Visan, M., *Global well-posedness and scattering for the defocusing energy-critical nonlinear Schrödinger equation in  $\mathbb{R}^{1+4}$* . Amer. J. Math. 129 (2007), 1–60.
- [28] Staffilani, G., *On the growth of high Sobolev norms of solutions for KdV and Schrödinger equations*, Duke Math. J., 86 (1997), 109-142.
- [29] Thirouin, J., work in preparation.
- [30] Xu, H., *Large time blow up for a perturbation of the cubic Szegő equation*, Anal. PDE, 7 (2014), No. 3, 717–731.
- [31] Zakharov, V. E., Shabat, A. B.: *Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media*. Soviet Physics JETP 34 (1972), no. 1, 62–69.
- [32] Zakharov, V., Guyenne, P., Pushkarev, A., Dias, F., *Wave turbulence in one-dimensional models*, Physica D 152–153 (2001) 573–619.

UNIVERSITÉ PARIS-SUD XI, LABORATOIRE DE MATHÉMATIQUES D’ORSAY, CNRS,  
 UMR 8628, ET INSTITUT UNIVERSITAIRE DE FRANCE  
*E-mail address:* Patrick.Gerard@math.u-psud.fr

FÉDÉRATION DENIS POISSON, MAPMO-UMR 6628, DÉPARTEMENT DE MATHÉMA-  
 TIQUES, UNIVERSITÉ D’ORLÉANS, 45067 ORLÉANS CEDEX 2, FRANCE  
*E-mail address:* Sandrine.Grellier@univ-orleans.fr